

CMPT 210: Probability and Computing

Lecture 22

Sharan Vaswani

November 26, 2024

Tail inequalities

- Variance gives us one way to measure how “spread” the distribution is.

Tail inequalities bound the probability that the r.v. takes a value much different from its mean.

Example: Consider a r.v. X that can take on only non-negative values and $\mathbb{E}[X] = 99.99$. Show that $\Pr[X \geq 300] \leq \frac{1}{3}$.

$$\begin{aligned} \text{Proof: } \mathbb{E}[X] &= \sum_{x \in \text{Range}(X)} x \Pr[X = x] = \sum_{x|x \geq 300} x \Pr[X = x] + \sum_{x|0 \leq x < 300} x \Pr[X = x] \\ &\geq \sum_{x|x \geq 300} (300) \Pr[X = x] + \sum_{x|0 \leq x < 300} x \Pr[X = x] \\ &= (300) \Pr[X \geq 300] + \sum_{x|0 \leq x < 300} x \Pr[X = x] \end{aligned}$$

If $\Pr[X \geq 300] > \frac{1}{3}$, then, $\mathbb{E}[X] > (300) \frac{1}{3} + \sum_{x|0 \leq x < 300} x \Pr[X = x] > 100$ (since the second term is always non-negative). Hence, if $\Pr[X \geq 300] > \frac{1}{3}$, $\mathbb{E}[X] > 100$ which is a contradiction since $\mathbb{E}[X] = 99.99$.

Markov's Theorem

Markov's theorem formalizes the intuition on the last slide and can be stated as follows.

Markov's Theorem: If X is a non-negative random variable, then for all $x > 0$,

$$\Pr[X \geq x] \leq \frac{\mathbb{E}[X]}{x}.$$

Proof: Define $\mathcal{I}\{X \geq x\}$ to be the indicator r.v. for the event $[X \geq x]$. Then for all values of X , $x\mathcal{I}\{X \geq x\} \leq X$.

$$\begin{aligned} \mathbb{E}[x\mathcal{I}\{X \geq x\}] \leq \mathbb{E}[X] &\implies x\mathbb{E}[\mathcal{I}\{X \geq x\}] \leq \mathbb{E}[X] \implies x\Pr[X \geq x] \leq \mathbb{E}[X] \\ &\implies \Pr[X \geq x] \leq \frac{\mathbb{E}[X]}{x}. \end{aligned}$$

Since the above theorem holds for all $x > 0$, we can set $x = c\mathbb{E}[X]$ for $c \geq 1$. In this case, $\Pr[X \geq c\mathbb{E}[X]] \leq \frac{1}{c}$. Hence, the probability that X is “far” from the mean in terms of the multiplicative factor c is upper-bounded by $\frac{1}{c}$.

Markov's Theorem – Example

Q: If X is a non-negative r.v. such that $\mathbb{E}[X] = 150$, bound the probability that X is at least 200.

Ans: $\Pr[X \geq 200] \leq \frac{\mathbb{E}[X]}{200} = \frac{3}{4}$

Q: If we are provided additional information that X can not take values less than 100 and $\mathbb{E}[X] = 150$, bound the probability that X is at least 200.

Define $Y := X - 100$. $\mathbb{E}[Y] = \mathbb{E}[X] - 100 = 50$ and Y is non-negative.

$$\Pr[X \geq 200] = \Pr[Y + 100 \geq 200] = \Pr[Y \geq 100] \leq \frac{\mathbb{E}[Y]}{100} = \frac{50}{100} = \frac{1}{2}$$

Hence, if we have additional information (in the form of a lower-bound that a r.v. can not be smaller than some constant $b > 0$), we can use Markov's Theorem on the shifted r.v. (Y in our example) and obtain a tighter bound on the probability of deviation.

Chebyshev's Theorem

Chebyshev's Theorem: For a r.v. X and any constant $y > 0$,

$$\Pr[|X - \mathbb{E}[X]| \geq y] \leq \frac{\text{Var}[X]}{y^2}.$$

Proof: Use Markov's Theorem with some cleverly chosen function of X . Formally, for some function f such that $Y := f(X)$ is non-negative. Using Markov's Theorem for Y ,

$$\Pr[f(X) \geq x] \leq \frac{\mathbb{E}[f(X)]}{x}$$

Choosing $f(X) = |X - \mathbb{E}[X]|^2$ and $x = y^2$ implies that $f(X)$ is non-negative and $x > 0$. Using Markov's Theorem,

$$\Pr[|X - \mathbb{E}[X]|^2 \geq y^2] \leq \frac{\mathbb{E}[|X - \mathbb{E}[X]|^2]}{y^2}$$

Note that $\Pr[|X - \mathbb{E}[X]|^2 \geq y^2] = \Pr[|X - \mathbb{E}[X]| \geq y]$, and hence,

$$\Pr[|X - \mathbb{E}[X]| \geq y] \leq \frac{\mathbb{E}[|X - \mathbb{E}[X]|^2]}{y^2} = \frac{\text{Var}[X]}{y^2}$$

Chebyshev's Theorem

- Chebyshev's Theorem bounds the probability that the random variable X is “far” away from the mean $\mathbb{E}[X]$ by an additive factor of x .
- If we set $x = c\sigma_X$ where σ_X is the standard deviation of X , then by Chebyshev's Theorem,

$$\Pr[(X \geq \mathbb{E}[X] + c\sigma_X) \cup (X \leq \mathbb{E}[X] - c\sigma_X)] = \Pr[|X - \mathbb{E}[X]| \geq c\sigma_X] \leq \frac{\text{Var}[X]}{c^2\sigma_X^2} = \frac{1}{c^2}$$

$$\Pr[\mathbb{E}[X] - c\sigma_X < X < \mathbb{E}[X] + c\sigma_X] = \Pr[|X - \mathbb{E}[X]| \leq c\sigma_X]$$

$$\implies \Pr[\mathbb{E}[X] - c\sigma_X < X < \mathbb{E}[X] + c\sigma_X] = 1 - \Pr[|X - \mathbb{E}[X]| \geq c\sigma_X] \geq 1 - \frac{1}{c^2}.$$

Hence, Chebyshev's Theorem can be used to bound the probability that X is “concentrated” near its mean.

- Unlike Markov's Theorem, Chebyshev's Theorem does not require the r.v. to be non-negative, but requires knowledge of the variance.

Chebyshev's Theorem - Example

Q: If X is a non-negative r.v. such that $\mathbb{E}[X] = 100$ and $\sigma_X = 15$, bound the probability that X is at least 300.

If we use Markov's Theorem, $\Pr[X \geq 300] \leq \frac{\mathbb{E}[X]}{300} = \frac{1}{3}$.


Note that $\Pr[|X - 100| \geq 200] = \Pr[X \leq -100 \cup X \geq 300] = \Pr[X \geq 300]$. Using Chebyshev's Theorem,

$$\Pr[X \geq 300] = \Pr[|X - 100| \geq 200] \leq \frac{\text{Var}[X]}{(200)^2} = \frac{15^2}{200^2} \approx \frac{1}{178}.$$

Hence, by exploiting the knowledge of the variance and using Chebyshev's inequality, we can obtain a tighter bound.

Chebyshev's Theorem - Example

Q: Consider a r.v. $X \sim \text{Bin}(20, 0.75)$. Plot the PDF_X , compute its mean and standard deviation and bound $\Pr[10 < X < 20]$.



$\text{Range}(X) = \{0, 1, \dots, 20\}$ and for $k \in \text{Range}(X)$,
 $f(k) = \binom{n}{k} p^k (1-p)^{n-k}$.

$$\mathbb{E}[X] = np = (20)(0.75) = 15$$

$\text{Var}[X] = np(1-p) = 20(0.75)(0.25) = 3.75$ and hence
 $\sigma_X = \sqrt{3.75} \approx 1.94$.

$$\begin{aligned} \Pr[10 < X < 20] &= 1 - \Pr[X \leq 10 \cup X \geq 20] \\ &= 1 - \Pr[|X - 15| \geq 5] \\ &= 1 - \Pr[|X - \mathbb{E}[X]| \geq 5] \\ &\geq 1 - \frac{\text{Var}[X]}{(5)^2} = 1 - \frac{3.75}{25} = 0.85. \end{aligned}$$

Hence, the “probability mass” of X is “concentrated” around its mean.